

---

## Technical Criteria for Reviewing Ancillary Services for Adults

---

### I. PHYSICAL THERAPY: REVIEW FOR BILLING AS ANCILLARY

- A. **STANDARDS OF PRACTICE:** The review process shall employ the standards of practice developed by the American Physical Therapy Association.
- B. Deficiency of function must be of a significant level that an ancillary clinician's expertise in designing or conducting a program in the presence of potential gain is documentable.
1. Therapeutic exercise
    - a. When exercising muscle or joint structure, the deficit requires a therapist's expertise to design, supervise, or conduct a program in which there is a need for functional or performance gain.
    - b. Progress is shown at predictable intervals.
    - c. Gradual progression is from passive to fully active range of motion per situation and reasonable goal.

#### Indication for Denial

- a. Lacks documented detail of dysfunction or goal.
- b. Goal seems unreasonable.
- c. Stability of resident questioned.
- d. Participation level a hindrance.
- e. Plateaued, goal achieved, or needs only repetitive range of motion for nursing care plan.
- f. Persistent flaccidity > 2-4 weeks in the focused area.

#### 2. Cold Therapy

- a. Pain or spasm reduction or adjustment to range of motion exercise (repeated cycles).
- b. Trigger point use myofascial pain syndrome.
- c. Spasticity.

#### Indication for Denial

- a. Response gain is not demonstrable.
- b. Performance is at nursing instructed level, and labile complex features.
- c. Inappropriate use in a vascular compromised setting (or labile or poor blood pressure control).
- d. Cold sensitivity disorder.

---

## Technical Criteria for Reviewing Ancillary Services for Adults

---

### 3. Low-Energy Laser

- a. Wound tissue healing.
- b. Pain management over trigger points.

#### **Indication for Denial**

- a. investigational.
- b. Effectiveness in rheumatoid arthritis questioned.

### 4. Transcutaneous Electric Nerve Stimulation (TENS)

- a. Post-operative incision pain.
- b. Orthopedic analgesia acute or chronic, application to either trigger point or peripheral nerve.
- c. Chronic low back pain...
- d. Osteogenesis.
- e. Reflex sympathetic dystrophy (RSD).

#### **Indication for Denial**

- a. Chronic radiculopathy pain.
- b. Cognitively impaired or unwilling to participate with schedule and safety factors.
- c. Unsafe application.
- d. Nursing is capable of managing (or resident can set-up, apply or control) after the initial evaluation of response or control setting is achieved.

### 5. Heat-Therapy

- a. Active treatment of musculoskeletal mobility or pain problem as part of a therapist-driven treatment plan.
- b. In conjunction with an exercise regimen.

#### **Indication for Denial**

- a. The active disorder is controlled, mostly for comfort.
  - b. Complexity manageable by nursing.
  - c. Resident is not responsive or is non-communicative.
  - d. Ischemic limbs or other site or atrophic skin.
-

---

## Technical Criteria for Reviewing Ancillary Services for Adults

---

### 6. Ultrasound

- a. Joint contracture or scar tissue before friction massage, stretch, or range of motion (ROM) exercise (intensities and durations still need work), i.e., post hip open reduction internal fixation.
- b. Reduce pain or muscle spasm.
- c. Trigger points.

#### **Indication for Denial**

- a. Use in precautionary situations.
- b. Impaired sensitivity or ischemia.
- c. Questionable efficacy such as chronic herpes zoster, hemiplegic shoulder pain, fresh wound, or chronic pressure sore.

### 7. Hydrotherapy

- a. Facilitate assistive or resistive exercise.
- b. Removal of exudated or necrotic tissue.
- c. Reduce muscle spasm or pain.

#### **Indication for Denial**

- a. General heat precautions.
- b. Treatment exposure using > 37 degrees centigrade in vascular impaired site.
- c. Absence of untoward effects or stable temperature tolerance and can be done by nursing staff.

### 8. Iontophoresis

- a. Antibiotic institution to avascular tissue.
- b. Medication for persistent post-surgical incision pain.
- c. Reduce inflammation or edema of musculoskeletal joints).

#### **Indication for Denial**

- a. Anesthetic use (injection faster).
  - b. Response lacking after reasonable interval.
-

---

## Technical Criteria for Reviewing Ancillary Services for Adults

---

### 9. Prosthesis

- a. Candidate has the capacity to use device.
- b. Candidate shows muscular strength, motor control, and range of motion adequate for gainful use.

#### Indication for Denial

- a. Unteachable.
- b. Lacks items in 9-a and b.
- c. Poor wound healing.
- d. Other inappropriate conditions (such as bilateral, above-knee amputation over age 45, or below-elbow amputee or flail joint shoulder or elbow).
- e. Repetitive exercises that nursing care plan can accomplish pre-prosthesis for stump shrinker use or prosthetic fitting.
- f. Repetitive use for distance or endurance only with level change having been achieved.
- g. Assisting routine care of equipment.
- h. Safety has been established so that the resident can perform trained exercise with supervision by nursing being the only need.

### 10. Electromyography Biofeedback

- a. Spasticity or weakness as part of an acute cerebral vascular accident (CVA).
- b. Acute or chronic spinal cord injury.
- c. Multiple sclerosis with mild spasticity.

#### Indication for Denial

- a. Absence of reasonable gain in the treatment plan time frame.
- b. Questionable effectiveness for the condition.
- c. Resident lacks voluntary control or motivation.

---

## Technical criteria for Reviewing Ancillary Services for Adults

---

### 11. High Pressure Wound Irrigation

- a. Heavily contaminated wound.

#### **Indication for Denial**

- a. Clean proliferating wounds
- b. Equipment or devices of questionable effectiveness or superiority to simpler devices.
- c. Nursing can provide equivalent service.

### 12. Hyperbaric Oxygen Wound Care

- a. Infected wounds or decubitus.
- b. Has reasonable circulation.

#### **Indication for Denial**

- a. Advanced ischemic area.
- b. Potential for thromboembolism.
- c. Severe vasospasm.
- d. Lack of significant improvement in 4 weeks.

---

## Technical criteria for Reviewing Ancillary Services for Adults

---

### II. OCCUPATIONAL THERAPY: REVIEW FOR BILLING AS ANCILLARY

A. - **STANDARDS OF PRACTICE:** The review process shall employ the standards of practice developed by the American Occupational Therapy Association..

B. Deficiency of function must be of a significant level that an ancillary-clinician's expertise in designing or conducting a program in the presence of potential gain is documentable.

#### 1. Therapeutic exercise

- a. When exercising muscle or joint structure the deficit requires a therapist's expertise to design, supervise, or conduct a program in which there is a need for functional or performance gain;
- b. Progress is shown at predictable intervals.
- c. Gradual progression is from passive to fully active range of motion per situation and reasonable goal.

#### **Indication for Denial**

- a. Lacks documented detail of dysfunction or goal.
- b. Goal seems unreasonable.
- c. Stability of the resident questioned.
- d.. Participation level is a hindrance.
- e. Plateaued, goal achieved, or needs only repetitive ROM for nursing care plan
- f. Persistent flaccidity > 2-4 weeks focused area.

#### 2. Shared Modalities for Physical Therapy

- a. Heat therapy.
- b.. Cold therapy.
- c. Prosthesis.
- d. Electromyography biofeedback.

. **Indication for Denial** (see listings for Physical Therapy)

#### 3. Functional Activities of Daily Living

- a. Feed.
- b. Dress.
- c. Bathe.
- d. Toileting.
- e. Grooming.

---

## Technical Criteria for Reviewing Ancillary Services for Adults

---

f. Cognition.

**. Indication for Denial**

- a. The condition prevents the individual from engaging in the technique or use of the device.
- b. Technique is reached, resident or nursing staff can maintain activities for endurance, distance or repetition.
- c. Chronic condition, therefore potential useful gain is questioned or minimal.
- d. Unable to advance or use more complex dexterity level due to cognitive limits.
- e. Biofeedback use in the presence of a prominent disorder speech, language use, cognition or volitional ability (inability to follow gestural or verbal instruction.)
- f. Coma stimulation - effectiveness questionable

---

## Technical Criteria for Reviewing Ancillary Services for Adults

---

### III. SPEECH THERAPY: REVIEW FOR BILLING AS ANCILLARY

- A. **STANDARDS OF PRACTICE:** The review process will employ the preferred practice patterns developed by the American Speech-Language-Hearing Association.

8; Deficiency of function must be of a significant level that an ancillary clinicians expertise in designing or conducting a program in the presence of potential gain is documentable.

#### 1. Treatment of Dysphagia (swallowing) Disorders

- a. Applicable diagnostic tests with confirmed abnormality (initial or progress recheck).
- b. Active teaching is appropriate for cognitive level (vs. delay till progress gain and provides alternative nutrition source).
- c. Uses specific postural, reflex facilitation, food placement, and modified diet techniques with demonstrable progress
- d. Prosthetic use

#### Indication for Denial

- a. Plateau, learned response, and repetitive exercise, reminders or prosthetics can be done by nursing as effectively.
- b. Confirmatory diagnostic test unavailable.
- c. Resident uncooperative or unreliable to safely use needed techniques.

#### 2. Speech and Cognitive Disorders

- a. Tentative projected rehabilitation goal at the stage when cognitive level permits measurable change.
- b. Participation by resident required for repetitive or grouped exercises.
- c. Prosthetic training.
- d. Demonstrates there is no contributing significant auditory impairment.
- e. Use of nursing facility environment or staff to assist goals.



---

## Technical Criteria for Reviewing Ancillary Services for Adults

---

### Indication for Denial

- a. Inability to participate.
- b. Plateau is reached in functional gain by measurable data or learned exercise and nursing can do repetitive technique.
- c. Effectiveness of modality or participation level is in question.
- d. Persisting active program beyond gain in condition having progressive deteriorating change or outlook (bilateral cerebral vascular accident, alzheimers).
- e. Oral-nonverbal apraxia beyond 2 months.
- f. Accompanying peripheral vision or hearing defects.

---

Technical Criteria for Reviewing Ancillary Services for Adults

---

**IV. OXYGEN THERAPY: REVIEW FOR MEDICAL NECESSITY**

- A. **STANDARDS OF PRACTICE:** The review process shall employ the Guidelines for Respiratory Care Services and Skilled Nursing Facilities developed jointly by the American Association of Respiratory Care and the American Health Care Association.
- B. Technical abbreviations-used in Item VII - Oxygen Therapy.
- .ABG - Arterial Blood Gases
  - AVF - Augmented Voltage Foot
  - O<sub>2</sub> - Oxygen Level
  - paO<sub>2</sub>- Partial Pressure of Oxygen
  - paCO<sub>2</sub> - Partial Pressure of Carbon Dioxide
  - Oxygen Stats - Oxygen Saturation Levels
  - HCT - Hematocrit Level
  - mm Hg - Millimeters of Mercury
- C. General Indicators.
1. PaO<sub>2</sub> < 55 mm Hg or saturation < 88% while breathing ambient air.
  2. Optimum medical management.
    - a. Ancillary respiratory medications.
    - b. Physiotherapy.
    - c. Associated adverse conditions addressed.
  3. PaO<sub>2</sub> of 56-59 mm Hg or saturation of 91% in the presence of one or more of the following:
    - a. Cor pulmonale (p wave greater than 3 mm in standard leads II, III, or AVF).
    - b. Right ventricular hypertrophy.
    - c. Erythrocytosis (Hct > 56%)
    - d. Reduced tissue oxygenation accompanied by neuropsych signs (i.e., tachycardia, tachypnea, dyspnea; cyanosis, diaphoresis chest pain or tightness, change in sensorium.
  4. For that resident whose clinical condition prohibits evaluation of arterial oxygen saturation without supplemental oxygen:
    - a. Oxygen saturation while on O<sub>2</sub> < 92%.
    - b. PaO<sub>2</sub> < 60 mm Hg.
-