

CABINET FOR HEALTH AND FAMILY SERVICES

Hospital Rate Improvement/UK/UL Directed Payment/MCO Value Based Payment Comparison

Medicaid Advisory Council

March 28, 2024



UK/UofL MEASURES

- Breast Cancer Screening
- Colorectal Cancer Screening
- Tobacco Use: Screening and Cessation Intervention
- Screening for Clinical Depression and Follow Up Plan
- ➤ Diabetes Care: Hemoglobin (HbA1c) Poor Control (>9%) (lower is better).
- Controlling High Blood Pressure (Hypertension)
- Statin Therapy for Patients with Cardiovascular Disease
- Body Mass Index (BMI) Screening and Follow-Up
- Medication Reconciliation Post-Discharge
- > 30 Day All Cause Readmissions (lower is better)
- Use of Opioids at High Dosage (lower is better)
- Childhood Immunization Status
- ➤ Well Child Visits, 3 6 years
- Well Child Visits, Rate 1: 0-15 Months
- Well Child Visits, Rate 2: 15-30 Months
- Screening for Depression and Follow-Up Plan: Ages 12 to 17
- Body Mass Index (BMI) Screening and Follow-Up: Ages 12 to 17
- Post-Partum Depression Screening
- SDOH Screening



HRIP MEASURES

- CAUTI Standardized Infection Ratio (SIR)
- U-Cath Ratio
- > CDI SIR
- > CDI Rate
- > Sepsis Screening
- > Sepsis Screening and Bundle Compliance
- Readmissions
- Concurrent e-Prescribing
- Opioid Education
- Admission Screening (HBIPS 1)
- ➤ Hours of Restraint (HBIPS 2)
- ➤ Hours of Seclusion (HBIPS 3)
- Rehab Discharge



VBP MEASURES

Core Measures

- ➤ HBD-Good Control (HbA1c<8)
- Child and Adolescent Well Care Visits 3-21 years of age (WCV) sum of stratifications (total)
- Childhood Immunization Status (CIS Combo 10)
- Immunization for Adolescents (IMA Combo 1)
- Postpartum Care (PPC) Postpartum Care
- Social Need Screening and Intervention (SNS-E)

Bonus Pool Measures

- Metabolic Monitoring for Children and Adolescents on Antipsychotic (APM-E)
- Follow-Up After Emergency Department Visit for Alcohol and Other Drug Dependence (FUA), Follow-Up Within 7 Days of ED Visit (Total)
- Weight Assessment and Counseling for Nutrition and Physical Activity for Children and Adolescents (WCC) Counseling for Nutrition Total
- Breast Cancer Screening (BCS-E)



UK/UgfL AND HRIP QVERLAP IN MEASURES

UK/UofL	HRIP
30 Day All Cause Readmissions (lower is better)	Readmissions
Use of Opioids at High Dosage (lower is better)	Opioid Education



UK/UofL AND VBP OVERLAP IN MEASURES

UK/UofL	VBP
Diabetes Care: Hemoglobin (HbA1c) Poor Control (>9%) (lower is better)	HBD-Good Control (HbA1c<8)
Well Child Visits, 3 – 6 years Well Child Visits, Rate 1: 0-15 Months Well Child Visits, Rate 2: 15-30 Months	Child and Adolescent Well Care Visits 3-21 years of age (WCV) sum of stratifications (total)
Post-Partum Depression Screening	Postpartum Care (PPC) Postpartum Care
SDOH Screening	Social Need Screening and Intervention (SNS-E)
Breast Cancer Screening	Breast Cancer Screening (BCS-E)

