



COVID-19 VACCINE AND BOOSTERS

Pfizer

Moderna

Johnson & Johnson (J&J)

Ages Recommended	5+ years old	18+ years old	18+ years old
Primary Series	2 doses Given 3 weeks (21 days) apart.	2 doses Given 4 weeks (28 days) apart.	1 dose
Booster Dose	You should get a booster dose at least 5 months after the last dose in their primary series.	You should get a booster dose at least 5 months after the last dose in their primary series.	You should get a booster dose of either Pfizer-BioNTech or Moderna (mRNA COVID-19 vaccines) at least 2 months after the first dose of J&J/Janssen COVID-19 Vaccine.
When am I "up-to-date"?	2 dose primary series less than 5 months ago OR booster 5 months after completion of primary series.	2 dose primary series less than 5 months ago OR booster 5 months after completion of primary series.	1 dose primary series less than 2 months ago OR booster 2 months after completion of primary series.
When am I "up-to-date" if I am moderately or severely immunocompromised?	2 dose series less than 28 days ago; or 1 additional dose (3rd dose to complete primary series) less than 3 months ago; or Booster (4th dose) 3 months after completion of primary 3 dose series.	2 dose series less than 28 days ago; or 1 additional dose (3rd dose to complete primary series) less than 3 months ago; or Booster (4th dose) 3 months after completion of primary 3 dose series.	1 dose primary series less than 28 days ago; 1 additional dose (2nd dose, Moderna or Pfizer) less than 2 months ago; Booster dose 2 months after second dose preferably with Moderna or Pfizer (3 total doses).
2nd Booster Dose	Ages 12+ immunocompromised Ages 50+	Ages 18+ immunocompromised Ages 50+	50+ or those that are immunocompromised SHOULD receive a second booster with a mRNA; 18-49 who received JJ for primary and booster MAY receive 2nd booster with mRNA