

Recommended Adult (with an additional risk factor or another indication) Immunization Schedule by Age Group, United States, 2022

Vaccine	19-26 Years	27-49 Years	50-64 Years	65+ Years
Influenza inactivated (IIV4) OR Influenza recombinant (RIV4)	1 dose annually	1 dose annually	1 dose annually	1 dose annually
Tetanus, diphtheria, pertussis (Tdap or Td)	1 dose Tdap each pregnancy; 1 dose Td/Tdap for wound management (talk with provider)	1 dose Tdap each pregnancy; 1 dose Td/Tdap for wound management (talk with provider)	1 dose Tdap each pregnancy; 1 dose Td/Tdap for wound management (talk with provider)	1 dose Tdap each pregnancy; 1 dose Td/Tdap for wound management (talk with provider)
Varicella (VAR)	2 doses (if born in 1980 or later)	2 doses (if born in 1980 or later)	2 doses	2 doses
Zoster recombinant (RZV)	2 doses for immunocompromising conditions (talk with provider)	2 doses for immunocompromising conditions (talk with provider)	2 doses	2 doses
Pneumococcal (PCV15, PCV20, PPSV23)	1 dose PCV15 followed by PPSV23 OR 1 dose PCV20 (talk with provider)	1 dose PCV15 followed by PPSV23 OR 1 dose PCV20 (talk with provider)	1 dose PCV15 followed by PPSV23 OR 1 dose PCV20 (talk with provider)	1 dose PCV15 followed by PPSV23 OR 1 dose PCV20 (talk with provider)
Hepatitis A (HepA)	2 or 3 doses depending on vaccine			
Hepatitis B (HepB)	2, 3, or 4 doses depending on vaccine or condition	2, 3, or 4 doses depending on vaccine or condition	2, 3, or 4 doses depending on vaccine or condition	2, 3, or 4 doses depending on vaccine or condition
Meningococcal A, C, W, Y (MenACWY)	1 or 2 doses depending on indication, see notes for booster recommendations	1 or 2 doses depending on indication, see notes for booster recommendations	1 or 2 doses depending on indication, see notes for booster recommendations	1 or 2 doses depending on indication, see notes for booster recommendations
Meningococcal B (MenB)	2 or 3 doses depending on vaccine and indication, see notes for booster recommendations	2 or 3 doses depending on vaccine and indication, see notes for booster recommendations	2 or 3 doses depending on vaccine and indication, see notes for booster recommendations	2 or 3 doses depending on vaccine and indication, see notes for booster recommendations
Haemophilus influenzae type b (Hib)	1 or 3 doses depending on indication			

COVID-19 vaccination is recommended for ages 6 months and older. A person is Up-to-Date with COVID-19 vaccinations when they have received all recommended COVID-19 vaccine doses, including any booster dose(s) when eligible.

