

Recommended Pediatric Immunization Schedule by Age Group, United States, 2022

Vaccine	at Birth (within 24 hours)	2 months	4 months	6 months	12 months	15 months	18 months	19-23 months	4-6 years	7-10 years	11-12 years	13-15 years	16-18 years
Hepatitis B (HepB)	1 dose	1 dose	1 dose (1)	1 dose	1 dose	1 dose	1 dose						
Diphtheria, tetanus, pertussis (DTaP/Tdap)		1 dose	1 dose	1 dose	1 dose (2)	1 dose (2)	1 dose (2)		1 dose		1 dose (Tdap)		
Haemophilus influenzae type b (Hib)		1 dose	1 dose	1 dose (1)	1 dose	1 dose							
Polio (IPV)		1 dose	1 dose	1 dose	1 dose	1 dose	1 dose		1 dose				
Pneumococcal conjugate (PCV13)		1 dose	1 dose	1 dose	1 dose	1 dose							
Rotavirus (RV)		1 dose	1 dose	1 dose (1)									
Measles, mumps, rubella (MMR)					1 dose	1 dose			1 dose				
Varicella (VAR)					1 dose	1 dose			1 dose				
Hepatitis A (HepA)					2 doses (6)	2 doses (6)	2 doses (6)	2 doses (6)	2 doses (6)	2 doses (6)	2 doses (6)	2 doses (6)	2 doses (6)
Human papillomavirus (HPV)											2 doses (3)		
Meningococcal A, C, W, Y (MenACWY)											1 dose		1 dose
Meningococcal B (MenB)													2 doses (4,5)
Influenza (Flu)				1 dose (7)	1 dose (7)	1 dose (7)	1 dose (7)	1 dose (7)	1 dose (7)	1 dose (7)	1 dose (7)	1 dose (7)	1 dose (7)

**COVID-19 vaccination is recommended for ages 6 months and older.
Talk to your child's healthcare provider about the vaccines recommended for their age.**

Footnotes:

1. Your child may not need this dose depending on the brand of vaccine that your healthcare provider uses.
2. This dose of DTaP may be given as early as age 12 months if it has been 6 months since the previous dose.
3. Children with certain medical conditions will need a third dose.
4. This vaccine may be given to healthy teens. It is also recommended for adolescents with certain health conditions.
5. Your teen may need an additional dose depending on your healthcare provider's recommendation.
6. 2 doses given 6 months apart routinely at age 12-23 months. HepA vaccine (2 doses) is also recommended for children and teens not previously vaccinated.
7. One dose each fall or winter to all people ages 6 months and older. Some children younger than age 9 years need 2 doses; ask your child's health-care provider if your child needs more than 1 dose.

