

Immunizations Update

Amy Herrington, DNP, RN, CEN, CNE

2023



Vaccination Media Competition

Submission deadline February 24th

Regional winners by March 3rd

State winner by March 10th



**POSTER & MEDIA
CONTEST**

Win
\$250

GRADES K THROUGH 12

BECOME A VACCINE CHAMPION

Illustrate the importance of routine childhood vaccines by designing a poster, brochure or short video.

Regional and state winners will be announced at the KHSAA Boys Basketball Tournament in March.

\$250 Prize sponsored by the Kentucky Association of Health Care Plans (KAHP) will be awarded to one state elementary, middle and high school student.

For official rules and additional information please visit:
<https://chfs.ky.gov/agencies/doh/deho/imm/BecomeAVaccineChampionRules.pdf>



**For Poster, Brochure & Video
Submission Details:**
Contact your Local Health Department

Let's Rise


Routine Immunizations on Schedule for Everyone (RISE)

[Print](#)



Let's RISE is a CDC initiative to provide actionable strategies, resources, and data to support getting all Americans back on-schedule with their routine immunizations to protect everyone from vaccine-preventable disease and disability.

<https://www.cdc.gov/vaccines/partners/routine-immunizations-lets-rise.html>



2023 Vaccine
Recommendations
2023

<https://www.cdc.gov/vaccines/acip/meetings/downloads/slides-2022-10-19-20/2023-immunization-schedule-508.pdf>

Table 1 Recommended Child and Adolescent Immunization Schedule for ages 18 years or younger, United States, 2023

These recommendations must be read with the notes that follow. For those who fall behind or start late, provide catch-up vaccination at the earliest opportunity as indicated by the green bars. To determine minimum intervals between doses, see the catch-up schedule (Table 2).

Vaccine	Birth	1 mo	2 mos	4 mos	6 mos	9 mos	12 mos	15 mos	18 mos	19–23 mos	2–3 yrs	4–6 yrs	7–10 yrs	11–12 yrs	13–15 yrs	16 yrs	17–18 yrs	
Hepatitis B (HepB)	1 st dose	← 2 nd dose →			← 3 rd dose →													
Rotavirus (RV): RV1 (2-dose series), RV5 (3-dose series)			1 st dose	2 nd dose	See Notes													
Diphtheria, tetanus, acellular pertussis (DTaP < 7 yrs)			1 st dose	2 nd dose	3 rd dose			← 4 th dose →				5 th dose						
Haemophilus influenzae type b (Hib)			1 st dose	2 nd dose	See Notes		← 3 rd or 4 th dose, See Notes →											
Pneumococcal conjugate (PCV13, PCV15)			1 st dose	2 nd dose	3 rd dose		← 4 th dose →											
Inactivated poliovirus (IPV < 18 yrs)			1 st dose	2 nd dose	← 3 rd dose →							4 th dose	See Notes					
COVID-19 (1vCOV-mRNA, 2vCOV-mRNA, 1vCOV-aPS)					2- or 3-dose primary series and booster (See Notes)													
Influenza (IV4) OR Influenza (LAIV4)					Annual vaccination 1 or 2 doses								OR	Annual vaccination 1 dose only				
Measles, mumps, rubella (MMR)					See Notes		← 1 st dose →					2 nd dose						
Varicella (VAR)							← 1 st dose →					2 nd dose						
Hepatitis A (HepA)					See Notes	2-dose series, See Notes												
Tetanus, diphtheria, acellular pertussis (Tdap ≥ 7 yrs)														1 dose				
Human papillomavirus (HPV)														See Notes				
Meningococcal (MenACWY-D ≥ 9 mos, MenACWY-CRM ≥ 2 mos, MenACWY-TT ≥ 2 years)			See Notes													1 st dose		2 nd dose
Meningococcal B (MenB-4C, MenB-FHbp)														See Notes				
Pneumococcal polysaccharide (PPSV23)												See Notes						
Dengue (DEN4CYD; 9–16 yrs)													Seropositive in endemic dengue areas (See Notes)					
<div><div></div>Range of recommended ages for all children</div> <div><div></div>Range of recommended ages for catch-up vaccination</div> <div><div></div>Range of recommended ages for certain high-risk groups</div> <div><div></div>Recommended vaccination can begin in this age group</div> <div><div></div>Recommended vaccination based on shared clinical decision-making</div> <div><div></div>No recommendation/not applicable</div>																		

Table 1 Recommended Child and Adolescent Immunization Schedule for ages 18 years or younger, United States, 2023

These recommendations must be read with the notes that follow. For those who fall behind or start late, provide catch-up vaccination at the earliest opportunity as indicated by the green bars. To determine minimum intervals between doses, see the catch-up schedule (Table 2).

Vaccine	Birth	1 mo	2 mos	4 mos	6 mos	9 mos	12 mos	15 mos	18 mos	19–23 mos	2–3 yrs	4–6 yrs	7–10 yrs	11–12 yrs	13–15 yrs	16 yrs	17–18 yrs
Hepatitis B (HepB)	1 st dose	← 2 nd dose →			← 3 rd dose →												
Rotavirus (RV): RV1 (2-dose series), RV5 (3-dose series)			1 st dose	2 nd dose	See Notes												
Diphtheria, tetanus, acellular pertussis (DTaP <7 yrs)			1 st dose	2 nd dose	3 rd dose			← 4 th dose →				5 th dose					
Haemophilus influenzae type b (Hib)			1 st dose	2 nd dose	See Notes		← 3 rd or 4 th dose → See Notes										
Pneumococcal conjugate (PCV13, PCV15)			1 st dose	2 nd dose	3 rd dose		← 4 th dose →										
Inactivated poliovirus (IPV <18 yrs)			1 st dose	2 nd dose	← 3 rd dose →							4 th dose					See Notes
COVID-19 (1vCOV-mRNA, 2vCOV-mRNA, 1vCOV-aPS)																	
Influenza (IIV4)																	
OR																	
Influenza (LAIV4)																	
Measles, mumps, rubella (MMR)					See Notes	← 1 st dose →						2 nd dose					
Varicella (VAR)						← 1 st dose →						2 nd dose					
Hepatitis A (HepA)					See Notes												
Tetanus, diphtheria, acellular pertussis (Tdap ≥7 yrs)															1 dose		
Human papillomavirus (HPV)															See Notes		
Meningococcal (MenACWY-D ≥9 mos, MenACWY-CRM ≥2 mos, MenACWY-TT ≥2 years)															1 st dose	2 nd dose	
Meningococcal B (MenB-4C, MenB-FHbp)																	
Pneumococcal polysaccharide (PPSV23)																	
Dengue (DEN4CYD; 9–16 yrs)																	

Range of recommended ages for all children
Range of recommended ages for catch-up vaccination
Range of recommended ages for certain high-risk groups
Recommended vaccination can begin in this age group
Recommended vaccination based on shared clinical decision-making
No recommendation/not applicable

Table 1

These recommendations must be read with the notes that follow. For those who fall behind or start late, provide catch-up vaccination at the earliest opportunity as indicated by the green bars. To determine minimum intervals between doses, see the catch-up schedule (Table 2).

Vaccine	Birth	1 mo	2 mos	4 mos	6 mos	9 mos	12 mos	15 mos	18 mos	19–23 mos	2–3 yrs	4–6 yrs	7–10 yrs	11–12 yrs	13–15 yrs	16 yrs	17–18 yrs				
Hepatitis B (HepB)	1 st dose	← 2 nd dose →			← 3 rd dose →																
Rotavirus (RV): RV1 (2-dose series), RV5 (3-dose series)			1 st dose	2 nd dose	See Notes																
Diphtheria, tetanus, acellular pertussis (DTaP <7 yrs)			1 st dose	2 nd dose	3 rd dose				← 4 th dose →			5 th dose									
Haemophilus influenzae type b (Hib)			1 st dose	2 nd dose	See Notes		← 3 rd or 4 th dose → See Notes														
Pneumococcal conjugate (PCV13, PCV15)			1 st dose	2 nd dose	3 rd dose		← 4 th dose →														
Inactivated poliovirus (IPV <18 yrs)			1 st dose	2 nd dose	← 3 rd dose →							4 th dose									
COVID-19 (1vCOV-mRNA, 2vCOV-mRNA, 1vCOV-aPS)	2- or 3-dose primary series and booster (See Notes)																				
Influenza (IV4) OR Influenza (LAIV4)					Annual vaccination 1 or 2 doses										Annual vaccination 1 dose only						
					Annual vaccination 1 or 2 doses										Annual vaccination 1 dose only						
Measles, mumps, rubella (MMR)					See Notes	← 1 st dose →					2 nd dose										
Varicella (VAR)						← 1 st dose →					2 nd dose										
Hepatitis A (HepA)					See Notes	2-dose series, See Notes															
Tetanus, diphtheria, acellular pertussis (Tdap ≥7 yrs)														1 dose							
Human papillomavirus (HPV)															See Notes						
Meningococcal (MenACWY-D ≥9 mos, MenACWY-CRM ≥2 mos, MenACWY-TT ≥2 years)			See Notes															1 st dose		2 nd dose	
Meningococcal B (MenB-4C, MenB-FHbp)														See Notes							
Pneumococcal polysaccharide (PPSV23)											See Notes										
Dengue (DEN4CYD; 9–16 yrs)													Seropositive in endemic dengue areas (See Notes)								

Range of recommended ages for all children

Range of recommended ages for catch-up vaccination

Range of recommended ages for certain high-risk groups

Recommended vaccination can begin in this age group

Recommended vaccination based on shared clinical decision-making

No recommendation/ not applicable

Table 2**Recommended Catch-up Immunization Schedule for Children and Adolescents Who Start Late or Who Are More than 1 Month Behind, United States, 2023**

The table below provides catch-up schedules and minimum intervals between doses for children whose vaccinations have been delayed. A vaccine series does not need to be restarted, regardless of the time that has elapsed between doses. Use the section appropriate for the child's age. **Always use this table in conjunction with Table 1 and the Notes that follow.**

Children age 4 months through 6 years					
Vaccine	Minimum Age for Dose 1	Minimum Interval Between Doses			
		Dose 1 to Dose 2	Dose 2 to Dose 3	Dose 3 to Dose 4	Dose 4 to Dose 5
Hepatitis B	Birth	4 weeks	8 weeks and at least 16 weeks after first dose minimum age for the final dose is 24 weeks		
Rotavirus	6 weeks Maximum age for first dose is 14 weeks, 6 days.	4 weeks	4 weeks maximum age for final dose is 8 months, 0 days		
Diphtheria, tetanus, and acellular pertussis	6 weeks	4 weeks	4 weeks	6 months	6 months
<i>Haemophilus influenzae</i> type b	6 weeks	No further doses needed if first dose was administered at age 15 months or older. 4 weeks if first dose was administered before the 1 st birthday. 8 weeks (as final dose) if first dose was administered at age 12 through 14 months.	No further doses needed if previous dose was administered at age 15 months or older. 4 weeks if current age is younger than 12 months and first dose was administered at younger than age 7 months and at least 1 previous dose was PRP-T (ActHib [®] , Pentacel [®] , Hibentac [®] , Vaxelis [®] or unknown) 8 weeks and age 12 through 59 months (as final dose) if current age is younger than 12 months and first dose was administered at age 7 through 11 months; OR if current age is 12 through 59 months and first dose was administered before the 1 st birthday and second dose was administered at younger than 15 months; OR if both doses were PedvaxHIB [®] and were administered before the 1 st birthday	8 weeks (as final dose) This dose only necessary for children age 12 through 59 months who received 3 doses before the 1 st birthday.	
Pneumococcal conjugate	6 weeks	No further doses needed for healthy children if first dose was administered at age 24 months or older. 4 weeks if first dose was administered before the 1 st birthday. 8 weeks (as final dose for healthy children) if first dose was administered at the 1 st birthday or after	No further doses needed for healthy children if previous dose was administered at age 24 months or older. 4 weeks if current age is younger than 12 months and previous dose was administered at <7 months old 8 weeks (as final dose for healthy children) if previous dose was administered between 7–11 months (wait until at least 12 months old); OR if current age is 12 months or older and at least 1 dose was administered before age 12 months	8 weeks (as final dose) this dose is only necessary for children aged 12 through 59 months regardless of risk, or age 60 through 71 months with any risk, who received 3 doses before age 12 months.	
Inactivated poliovirus	6 weeks	4 weeks	4 weeks if current age is <4 years 6 months (as final dose) if current age is 4 years or older	6 months (minimum age 4 years for final dose)	
Measles, mumps, rubella	12 months	4 weeks			
Varicella	12 months	3 months			
Hepatitis A	12 months	6 months			
Meningococcal ACWY	2 months MenACWY-CRM 9 months MenACWY-D 2 years MenACWY-TT	8 weeks	See Notes	See Notes	
Children and adolescents age 7 through 18 years					
Meningococcal ACWY	Not applicable (N/A)	8 weeks			
Tetanus, diphtheria, tetanus, diphtheria, and acellular pertussis	7 years	4 weeks	4 weeks if first dose of DTaP/DT was administered before the 1 st birthday 6 months (as final dose) if first dose of DTaP/DT or Tdap/Td was administered at or after the 1 st birthday	6 months if first dose of DTaP/DT was administered before the 1 st birthday	
Human papillomavirus	9 years	Routine dosing intervals are recommended.			
Hepatitis A	N/A	6 months			
Hepatitis B	N/A	4 weeks	8 weeks and at least 16 weeks after first dose		
Inactivated poliovirus	N/A	4 weeks	6 months A fourth dose is not necessary if the third dose was administered at age 4 years or older and at least 6 months after the previous dose.	A fourth dose of IPV is indicated if all previous doses were administered at <4 years or if the third dose was administered <6 months after the second dose.	
Measles, mumps, rubella	N/A	4 weeks			
Varicella	N/A	3 months if younger than age 13 years. 4 weeks if age 13 years or older			
Dengue	9 years	6 months	6 months		

Notes

Recommended Child and Adolescent Immunization Schedule for ages 18 years or younger, United States, 2023

For vaccination recommendations for persons ages 19 years or older, see the Recommended Adult Immunization Schedule, 2023.

Additional information

- * Consult relevant ACIP statements for detailed recommendations at www.cdc.gov/vaccines/hcp/acip-recs/index.html.
- * For calculating intervals between doses, 4 weeks = 28 days. Intervals of ≥ 4 months are determined by calendar months.
- * Within a number range (e.g., 12–18), a dash (–) should be read “and”.
- * Vaccine age or minimum age for administration should be read “at least”.
- * For vaccination of persons with immunodeficiencies, see Table 8-1, Vaccination of persons with primary and secondary immunodeficiencies, in *General Best Practice Guidelines for Immunization* at www.cdc.gov/vaccines/hcp/acip-recs/general-recs/immunocompetence.html, and Immunization

COVID-19 vaccination
(minimum age: 6 months [Moderna and Pfizer-BioNTech COVID-19 vaccines], 12 years [Novavax COVID-19 Vaccine])

Routine vaccination

- * **Primary series:**
 - Age 6 months–4 years: 2-dose series at 0, 4–8 weeks (Moderna) or 3-dose series at 0, 3–8, 11–16 weeks (Pfizer-BioNTech)
 - Age 5–11 years: 2-dose series at 0, 4–8 weeks (Moderna) or 3-dose series at 0, 3–8, 11–16 weeks (Pfizer-BioNTech)

Routine vaccination

- Add bullet: **Dengue vaccine should not be administered to children traveling to or visiting endemic dengue areas.**

Dengue vaccination
(minimum age: 9 years)

Routine vaccination

- * Age 9–16 years living in areas with endemic dengue **AND** have laboratory confirmation of previous dengue infection
 - 3-dose series administered at 0, 6, and 12 months
- * Endemic areas include Puerto Rico, American Samoa, US Virgin Islands, Federated States of Micronesia, Republic of Marshall Islands, and the Republic of Palau. For updated guidance on dengue endemic areas and pre-vaccination laboratory testing see www.cdc.gov/mmwr/volumes/70/rr/rr7006a1.htm?s_cid=rr7006a1_w and www.cdc.gov/dengue/vaccine/hcp/index.html
- * Dengue vaccine should not be administered to children traveling to or visiting endemic dengue areas.

Diphtheria, tetanus, and pertussis (DTaP) vaccination (minimum age: 6 weeks [4 years for Kinrix® or Quadracel®])

Routine vaccination

- * 5-dose series at age 2, 4, 6, 15–18 months, 4–6 years
 - **Prospectively:** Dose 4 may be administered as early as age 12 months if at least 6 months have elapsed since dose 3.
 - **Retrospectively:** A 4th dose that was inadvertently administered as early as age 12 months may be counted if at least 4 months have elapsed since dose 3.

Catch-up vaccination

- * Dose 5 is not necessary if dose 4 was administered at age

* **Adults** (persons aged 19 years or older): 4-dose series (2-dose primary series at age 2, 4, and 6 months, followed by a booster dose* at age 12–15 months)

- *Vaxelis[®] is not recommended for use as a booster dose. A different Hib-containing vaccine should be used for the booster dose.

* **PedvaxHIB[®]**: 3-dose series (2-dose primary series at age 2 and 4 months, followed by a booster dose at age 12–15 months)

SICKLE CELL DISEASE:

Age 12–59 months

- Unvaccinated or only 1 dose before age 12 months:

2 doses, 8 weeks apart

- 2 or more doses before age 12 months:

1 dose at least 8 weeks after previous dose

Unvaccinated* persons age 5 years or older

- 1 dose

* Fluorine-containing vaccine

HEPATITIS B VACCINE

• Persons traveling to or working in countries with high or intermediate endemic hepatitis A (www.cdc.gov/travel/):

- **Infants age 6–11 months:** 1 dose before departure; revaccinate with 2 doses (separated by at least 6 months) between age 12–23 months.

- **Unvaccinated age 12 months or older:** Administer dose 1 as soon as travel is considered.

Hepatitis B vaccination (minimum age: birth)

Routine vaccination

• 3-dose series at age 0, 1–2, 6–18 months (use **monovalent HepB vaccine for doses administered before age 6 weeks**)

- Birth weight $\geq 2,000$ grams: 1 dose within 24 hours of birth if medically stable

- Birth weight $< 2,000$ grams: 1 dose at chronological age 1 month or hospital discharge (whichever is earlier and even if weight is still $< 2,000$ grams).

• Infants who did not receive a birth dose should begin the series as soon as possible (see Table 2 for minimum intervals).

• Administration of 4 doses is permitted when a combination vaccine containing HepB is used after the birth dose.

• **Minimum intervals** (see Table 2): when 4 doses are administered, substitute “dose 4” for “dose 3” in these calculations

• **Final (3rd or 4th) dose:** age 6–18 months (minimum age 24 weeks)

• **Mother is HBsAg-positive**

- **Birth dose (monovalent HepB vaccine only):** administer HepB vaccine and hepatitis B immune globulin (HBIG) (in separate limbs) within 12 hours of birth, regardless of birth weight.

- **Birth weight $< 2,000$ grams:** administer 3 additional doses of HepB vaccine beginning at age 1 month (total of 4 doses)

- **Final (3rd or 4th) dose:** administer at age 6 months (minimum age 24 weeks)

- **Test for HBsAg and anti-HBs at age 9–12 months.** If HepB series is delayed, test 1–2 months after final dose. Do not test before age 9 months.

Routine vaccination

Mother is HBsAg-positive

- **Birth dose (monovalent HepB vaccine only):** administer HepB vaccine and hepatitis B immune globulin (HBIG) (in separate limbs) within 12 hours of birth, regardless of birth weight.

- **Birth weight $< 2,000$ grams:** administer 3 additional doses of HepB vaccine beginning at age 1 month (total of 4 doses)

- **Final (3rd or 4th) dose:** administer at age 6 months (minimum age 24 weeks)

- **Test for HBsAg and anti-HBs at age 9–12 months.** If HepB series is delayed, test 1–2 months. Do not test before age 9 months.

Doses administered within 14 days of starting therapy or receiving therapy should be repeated at least 3 months

older should receive dose 2 at least 6 months after dose 1.

Notes

Recommended Child and Adolescent Immunization Schedule

• **Mother is HBsAg-unknown**

If other evidence suggestive of maternal hepatitis B infection exists (e.g., presence of HBV DNA, HBeAg-positive, or mother known to have chronic hepatitis B infection), manage infant as if mother is HBsAg-positive

- **Birth dose (monovalent HepB vaccine only):**

• Birth weight $\geq 2,000$ grams: administer **HepB vaccine** within 12 hours of birth. Determine mother's HBsAg status as soon as possible. If mother is determined to be HBsAg-positive, administer **HBIG** as soon as possible (in separate limb), but no later than 7 days of age.

• Birth weight $< 2,000$ grams: administer **HepB vaccine** and **HBIG** (in separate limbs) within 12 hours of birth. Administer 3 additional doses of **HepB vaccine** beginning at age 1 month (total of 4 doses)

- **Final (3rd or 4th) dose:** administer at age 6 months (minimum age 24 weeks)

- If mother is determined to be HBsAg-positive or if status remains unknown, test for HBsAg and anti-HBs at age 9–12 months. If HepB series is delayed, test 1–2 months after final dose. Do not test before age 9 months.

Catch-up vaccination

• Unvaccinated persons should complete a 3-dose series at 0, 1–2, 6 months. See Table 2 for minimum intervals

• Adolescents age 11–15 years may use an alternative 2-dose schedule with at least 4 months between doses (adult formulation **Recombivax HB**® only).

• Adolescents age 18 years or older may receive:

- **HepBisav-B**®: 2-dose series at least 4 weeks apart

- **PreHevbrio**®: 3-dose series at 0, 1, and 6 months

- Combined HepA and HepB vaccine, **Twinrix**®: 3-dose series (0, 1, and 6 months) or 4-dose series (3 doses at 0, 7, and 21–30 days, followed by a booster dose at 12 months).

Special situations

• Revaccination is not generally recommended for persons with a normal immune status who were vaccinated as infants, children, adolescents, or adults.

• Post-vaccination serology testing and revaccination (if anti-HBs < 10 mIU/mL) is recommended for certain populations, including:

- Infants born to HBsAg-positive mothers

- Persons who are predialysis or on maintenance dialysis

- Other immunocompromised persons

- For detailed revaccination recommendations, see www.cdc.gov/vaccines/hcp/acip-recs/vacc-specific/hepb.html.

Note: HepBisav-B and PreHevbrio are not recommended in pregnancy due to lack of safety data in pregnant women

Hepatitis B vaccination

Routine vaccination

Mother is HBsAg-unknown

If other evidence suggestive of maternal hepatitis B infection exists (e.g., presence of HBV DNA, HBeAg-positive, or mother known to have chronic hepatitis B infection), manage infant as if mother is HBsAg-positive

- **Birth dose (monovalent HepB vaccine only):**

• Birth weight $\geq 2,000$ grams: administer **HepB vaccine** within 12 hours of birth. Determine mother's HBsAg status as soon as possible. If mother is determined to be HBsAg-positive, administer **HBIG** as soon as possible (in separate limb), but no later than 7 days of age.

• Birth weight $< 2,000$ grams: administer **HepB vaccine** and **HBIG** (in separate limbs) within 12 hours of birth. Administer 3 additional doses of **HepB vaccine** beginning at age 1 month (total of 4 doses)

- **Final (3rd or 4th) dose:** administer at age 6 months (minimum age 24 weeks)

- If mother is determined to be HBsAg-positive or if status remains unknown, test for HBsAg and anti-HBs at age 9–12 months. If HepB series is delayed, test 1–2 months after final dose. Do not test before age 9 months.

Human papillomavirus (HPV) vaccine (minimum age 9 years)

Routine vaccination

• HPV vaccine (can start as early as age 9)

• Recommended schedule

• 2-dose series

• 1 dose for all persons age 9 years or older

• 2 doses, 6–12 months apart

• 1 dose for all persons age 9 years or older

• 2 doses, 6–12 months apart

• 1 dose for all persons age 9 years or older

• 2 doses, 6–12 months apart

• 1 dose for all persons age 9 years or older

• 2 doses, 6–12 months apart

• 1 dose for all persons age 9 years or older

• 2 doses, 6–12 months apart

• 1 dose for all persons age 9 years or older

• 2 doses, 6–12 months apart

• 1 dose for all persons age 9 years or older

• 2 doses, 6–12 months apart

• 1 dose for all persons age 9 years or older

• 2 doses, 6–12 months apart

• 1 dose for all persons age 9 years or older

• 2 doses, 6–12 months apart

• 1 dose for all persons age 9 years or older

• 2 doses, 6–12 months apart

• 1 dose for all persons age 9 years or older

• 2 doses, 6–12 months apart

• 1 dose for all persons age 9 years or older

• 2 doses, 6–12 months apart

• 1 dose for all persons age 9 years or older

• 2 doses, 6–12 months apart

• 1 dose for all persons age 9 years or older

• 2 doses, 6–12 months apart

• 1 dose for all persons age 9 years or older

• 2 doses, 6–12 months apart

• 1 dose for all persons age 9 years or older

• 2 doses, 6–12 months apart

• 1 dose for all persons age 9 years or older

• 2 doses, 6–12 months apart

• 1 dose for all persons age 9 years or older

• 2 doses, 6–12 months apart

• 1 dose for all persons age 9 years or older

• 2 doses, 6–12 months apart

• 1 dose for all persons age 9 years or older

• 2 doses, 6–12 months apart

• 1 dose for all persons age 9 years or older

Notes

Recommended Child and Adolescent Immunization Schedule for ages 18 years or younger, United States, 2023

• **Mother is HBsAg-unknown**

If other evidence suggestive of maternal hepatitis B infection exists (e.g., presence of HBV DNA, HBeAg-positive, or mother known to have chronic hepatitis B infection), manage infant as if mother is HBsAg-positive

- **Birth dose (monovalent HepB vaccine only):**

Birth weight $\geq 2,000$ grams: administer HepB vaccine within 12 hours of birth. Determine mother's HBsAg status as soon as possible. If mother is determined to be HBsAg-positive, administer HBIG as soon as possible (in separate limb), but no later than 7 days of age.

Birth weight $< 2,000$ grams: administer HepB vaccine and HBIG (in separate limbs) within 12 hours of birth. Administer 3 additional doses of HepB vaccine beginning at age 1 month (total of 4 doses)

- **Final (3rd or 4th) dose:** administer at age 6 months (minimum age 24 weeks)

- If mother is determined to be HBsAg-positive or if status remains unknown, test for HBsAg and anti-HBs at age 9–12 months. If HepB series is delayed, test 1–2 months after final dose. Do not test before age 9 months.

Catch-up vaccination

• Unvaccinated persons should complete a 3-dose series at 0, 1–2, 6 months. See Table 2 for minimum intervals

• Adolescents age 11–15 years may use an alternative 2-dose schedule with at least 4 months between doses (adult formulation **Recombivax HB**® only).

• Adolescents age 18 years or older may receive:

- **HepBisav-B**®: 2-dose series at least 4 weeks apart

- **PreHevbrio**®: 3-dose series at 0, 1, and 6 months

- Combined HepA and HepB vaccine, **Twinrix**®: 3-dose series (0, 1, and 6 months) or 4-dose series (3 doses at 0, 7, and 21–30 days, followed by a booster dose at 12 months).

Special situations

• Revaccination is not generally recommended for persons with a normal immune status who were vaccinated as infants, children, adolescents, or adults.

• **Post-vaccination serology testing and revaccination** (if anti-HBs < 10 mIU/mL) is recommended for certain populations, including:

- Infants born to HBsAg-positive mothers

- Persons who are predialysis or on maintenance dialysis

- Other immunocompromised persons

- For detailed revaccination recommendations, see www.cdc.gov/vaccines/hcp/acip-recs/vacc-specific/hepb.html.

Note: HepBisav-B and PreHevbrio are not recommended in pregnancy due to lack of safety data in pregnant women

Human papillomavirus vaccination (minimum age: 9 years)

Routine and catch-up vaccination

• HPV vaccination routinely recommended at age 11–12 years (can start at age 9 years) and catch-up HPV vaccination recommended for all persons through age 18 years if not adequately vaccinated

• 2- or 3-dose series depending on age at initial vaccination:

- Age 9–14 years at initial vaccination: 2-dose series at 0, 6–12 months (minimum interval: 5 months; repeat dose if administered too soon)

- Age 15 years or older at initial vaccination: 3-dose series at 0, 1–2, 3–5 months

• Interrupted series: If a series is interrupted, restart at the next dose

• No additional doses are recommended for persons who have received all recommended doses

• For the 2022–2023 season, see www.cdc.gov/mmwr/volumes/71/mr7101a1.html.

• For the 2023–24 season, see the 2023–24 ACIP influenza vaccine recommendations.

• **Special situations**

• Egg allergy, hives only: Any influenza vaccine appropriate for age and health status annually

• Egg allergy with symptoms other than hives (e.g., angioedema, respiratory distress) or required epinephrine or another emergency medical intervention: Any influenza vaccine appropriate for age and health status may be administered. If using egg-based IV4 or LAIV4, administer

• For the 2022–2023 season, see www.cdc.gov/mmwr/volumes/71/mr7101a1.html.

• For the 2023–24 season, see the 2023–24 ACIP influenza vaccine recommendations.

• **Special situations**

• Egg allergy, hives only: Any influenza vaccine appropriate for age and health status annually

• Egg allergy with symptoms other than hives (e.g., angioedema, respiratory distress) or required epinephrine or another emergency medical intervention: Any influenza vaccine appropriate for age and health status may be administered. If using egg-based IV4 or LAIV4, administer

• For the 2022–2023 season, see www.cdc.gov/mmwr/volumes/71/mr7101a1.html.

• For the 2023–24 season, see the 2023–24 ACIP influenza vaccine recommendations.

Hepatitis B vaccination

Catch-up vaccination

Added bullet:

- Adolescents aged 18 years or older may receive:
 - **HepBisav-B**®: 2-dose series at least 4 weeks apart
 - **PreHevbrio**®: 3-dose series at 0, 1, and 6 months
 - Combined HepA and HepB vaccine, **Twinrix**®: 3-dose series (0, 1, and 6 months) or 4-dose series (3 doses at 0, 7, and 21–30 days, followed by a booster dose at 12 months).

Special situations

Added a sentence:

* Menveo has two formulations: One-vial (all liquid) and Two-vial (lyophilized and liquid). Menveo one-vial formulation should **NOT** be used before age 10 years.

Meningococcal serogroup A,C,W,Y vaccination (minimum age: 2 months [MenACWY-CRM, Menveo], 9 months [MenACWY-D, Menactra], 2 years [MenACWY-TT, MenQuadfi])

Routine vaccination

- 2-dose series at age 11–12 years; 16 years

Catch-up vaccination

- Age 13–15 years: 1 dose now and booster at age 16–18 years (minimum interval: 8 weeks)
- Age 16–18 years: 1 dose

Special situations

Anatomic or functional asplenia (including sickle cell disease), HIV infection, persistent complement component deficiency, complement inhibitor (e.g., eculizumab, ravulizumab) use:

- **Menveo****
 - Dose 1 at age 2 months: 4-dose series (additional 3 doses at age 4, 6 and 12 months)
 - Dose 1 at age 3–6 months: 3- or 4- dose series (dose 2 [and dose 3 if applicable] at least 8 weeks after previous dose until a dose is received at age 7 months or older, followed by an additional dose at least 12 weeks later and after age 12 months)
 - Dose 1 at age 7–23 months: 2-dose series (dose 2 at least 12 weeks after dose 1 and after age 12 months)
 - Dose 1 at age 24 months or older: 2-dose series: at least 8 weeks apart
- **Menactra***
 - **Persistent complement component deficiency or complement inhibitor use:**
 - Age 9–23 months: 2-dose series at least 12 weeks apart
 - Age 24 months or older: 2-dose series at least 8 weeks apart

Travel in countries with hyperendemic or epidemic meningococcal disease, including countries in the African meningitis belt or during the Hajj (www.cdc.gov/travel/):

- Children less than age 24 months:
 - **Menveo** (age 2–23 months)**
 - Dose 1 at age 2 months: 4-dose series (additional 3 doses at age 4, 6 and 12 months)
 - Dose 1 at age 3–6 months: 3- or 4- dose series (dose 2 [and dose 3 if applicable] at least 8 weeks after previous dose until a dose is received at age 7 months or older, followed by an additional dose at least 12 weeks later and after age 12 months)
 - Dose 1 at age 7–23 months: 2-dose series (dose 2 at least 12 weeks after dose 1 and after age 12 months)
 - **Menactra* (age 9–23 months)**
 - 2-dose series (dose 2 at least 12 weeks after dose 1; dose 2 may be administered as early as 8 weeks after dose 1 in travelers)

- Children age 2 years or older: 1 dose Menveo*, Menactra*, or MenQuadfi*

First-year college students who live in residential housing (if not previously vaccinated at age 16 years or older) or military recruits:

- 1 dose Menveo**, Menactra*, or MenQuadfi*

Adolescent vaccination of children who received MenACWY prior to age 10 years:

- **Children for whom boosters are recommended** because of an ongoing increased risk of meningococcal disease (e.g., those with complement deficiency, HIV, or asplenia): Follow the booster schedule for persons at increased risk.
- **Children for whom boosters are not recommended** (e.g., a healthy child who received a single dose for travel to a country where meningococcal disease is endemic): Administer MenACWY according to the recommended adolescent schedule with dose 1 at age 11–12 years and dose 2 at age 16 years.

for ages 18 years or younger, United States, 2023

*Menveo has two formulations: One-vial (all liquid) and Two-vial (lyophilized and liquid). Menveo one-vial formulation should **NOT** be used before age 10 years.

Note: Menactra* should be administered either before or at the same time as DTaP. MenACWY vaccines may be administered simultaneously with MenB vaccines if indicated, but at a different anatomic site, if feasible.

For MenACWY booster dose recommendations for groups listed under "Special situations" and in an outbreak setting and additional meningococcal vaccination information, see www.cdc.gov/mmwr/volumes/69/rr/r6909a1.htm.

Meningococcal serogroup B vaccination (minimum age: 10 years [MenB-4C, Bexsero*, MenB-FHbp, Trumenba*])

Shared clinical decision-making

- Adolescents not at increased risk age 16–23 years (preferred age 16–18 years) based on shared clinical decision-making:
 - Bexsero*: 2-dose series at least 1 month apart
 - Trumenba*: 2-dose series at least 6 months apart (if dose 2 is administered earlier than 6 months, administer a 3rd dose at least 4 months after dose 2)

Special situations

Anatomic or functional asplenia (including sickle cell disease), persistent complement component deficiency, complement inhibitor (e.g., eculizumab, ravulizumab) use:

- Bexsero*: 2-dose series at least 1 month apart
- Trumenba*: 3-dose series at 0, 1–2, 6 months (if dose 2 was administered at least 6 months after dose 1, dose 3 not needed; if dose 3 is administered earlier than 4 months after dose 2, a fourth dose should be administered at least 4 months after dose 3)

Note: Bexsero* and Trumenba* are not interchangeable; the same product should be used for all doses in a series.

For MenB booster dose recommendations for groups listed under "Special situations" and in an outbreak setting and additional meningococcal vaccination information, see www.cdc.gov/mmwr/volumes/69/rr/r6909a1.htm.

Notes

Recommended Child and Adolescent Immunization Schedule for ages 18 years

Pneumococcal vaccination (minimum age: 6 weeks [PCV13], [PCV15], 2 years [PPSV23])

Routine vaccination with PCV

- 4-dose series at 2, 4, 6, 12–15 months

Catch-up vaccination with PCV

- Healthy children age 24–59 months with any incomplete* PCV series: 1 dose PCV
- For other catch-up guidance, see Table 2.

Note: PCV13 and PCV15 can be used interchangeably for children who are healthy or have underlying conditions. No additional PCV15 is indicated for children who have received 4 doses of PCV13 or another age appropriate complete PCV13 series.

Special situations

Underlying conditions below: When both PCV and PPSV23 are indicated, administer PCV first. PCV and PPSV23 should not be administered during same visit.

Chronic heart disease (particularly cyanotic congenital heart disease and cardiac failure); **chronic lung disease** (including asthma treated with high-dose, oral corticosteroids); **diabetes mellitus**

Age 2–5 years

- Any incomplete* series with:
 - 3 PCV doses: 1 dose PCV (at least 8 weeks after any prior PCV dose)
 - Less than 3 PCV doses: 2 doses PCV (8 weeks after the most recent dose and administered 8 weeks apart)
- No history of PPSV23: 1 dose PPSV23 (at least 8 weeks after completing all recommended PCV doses)

Age 6–18 years

- Any incomplete* series with PCV: no further PCV doses needed
- No history of PPSV23: 1 dose PPSV23 (at least 8 weeks after completing all recommended PCV doses)

Cerebrospinal fluid leak, cochlear implant:

Age 2–5 years

- Any incomplete* series with:
 - 3 PCV doses: 1 dose PCV (at least 8 weeks after any prior PCV dose)
 - Less than 3 PCV doses: 2 doses PCV (8 weeks after the most recent dose and administered 8 weeks apart)
- No history of PPSV23: 1 dose PPSV23 (at least 8 weeks after any prior PCV dose)

Age 6–18 years

- No history of either PCV or PPSV23: 1 dose PCV, 1 dose PPSV23 at least 8 weeks later
- Any PCV but no PPSV23: 1 dose PPSV23 at least 8 weeks after the most recent dose of PCV
- PPSV23 but no PCV: 1 dose PCV at least 8 weeks after the most recent dose of PPSV23

Sickle cell disease and other hemoglobinopathies; anatomic or functional asplenia; congenital or acquired immunodeficiency; HIV infection; chronic renal failure; nephrotic syndrome; malignant neoplasms, leukemias, lymphomas, Hodgkin disease, and other diseases associated with treatment with immunosuppressive drugs or radiation therapy; solid organ transplantation; multiple myeloma:

Age 2–5 years

- Any incomplete* series with:
 - 3 PCV doses: 1 dose PCV (at least 8 weeks after any prior PCV dose)
 - Less than 3 PCV doses: 2 doses PCV (8 weeks after the most recent dose and administered 8 weeks apart)
- No history of PPSV23: 1 dose PPSV23 (at least 8 weeks after any prior PCV dose) and a dose 2 of PPSV23 5 years later

Age 6–18 years

- No history of either PCV or PPSV23: 1 dose PCV, 2 doses PPSV23 (dose 1 of PPSV23 administered 8 weeks after PCV and dose 2 of PPSV23 administered at least 5 years after dose 1 of PPSV23)
- Any PCV but no PPSV23: 2 doses PPSV23 (dose 1 of PPSV23 administered 8 weeks after the most recent dose of PCV and dose 2 of PPSV23 administered at least 5 years after dose 1 of PPSV23)
- PPSV23 but no PCV: 1 dose PCV at least 8 weeks after the most recent PPSV23 dose and a dose 2 of PPSV23 administered 5 years after dose 1 of PPSV23 and at least 8 weeks after a dose of PCV

*Incomplete series = Not having received all doses in either the recommended series or an age-appropriate catch-up series see Table 2 in ACIP pneumococcal recommendations at www.cdc.gov/mmwr/volumes/71/wr/mm7137a3.htm

Routine, Catch-up, and Special situations

- Added PCV15
- Replaced PCV13 with PCV
- Added note: PCV13 and PCV15 can be used interchangeably for children who are healthy or have underlying conditions. No additional PCV15 is indicated for children who have received 4 doses of PCV13 or another age appropriate complete PCV13 series.
- Deleted bullet: *Chronic liver disease, alcoholism*

Notes

Recommended Child and Adolescent Immunization Schedule for ages 18 years or younger, United States, 2023

Pneumococcal vaccination

(minimum age: 6 weeks [PCV13], [PCV15], 2 years [PPSV23])

Routine vaccination with PCV

• 4-dose series at 2, 4, 6, 12–15 months

Catch-up vaccination with PCV

• Healthy children age 24–59 months with any incomplete PCV series: 1 dose PCV

• For other catch-up guidance, see Table 2.

Note: PCV13 and PCV15 can be used interchangeably for children who are healthy or have underlying conditions. No additional PCV15 is indicated for children who have received 4 doses of PCV13 or another age-appropriate complete PCV13 series.

Special situations

Underlying conditions below: When both PCV and PPSV23 are indicated, administer PCV first. PCV and PPSV23 should not be administered during same visit.

Chronic heart disease (particularly cyanotic congenital heart disease and cardiac failure); chronic lung disease (including asthma treated with high-dose, oral corticosteroids); diabetes mellitus.

Age 2–5 years

• Any incomplete series with:
- 3 PCV doses: 1 dose PCV (at least 8 weeks after previous final dose)

Age 6–18 years

• No history of either PCV or PPSV23: 1 dose PCV, 1 dose PPSV23 at least 8 weeks later

• Any PCV but no PPSV23: 1 dose PPSV23 at least 8 weeks after the most recent dose of PCV

• PPSV23 but no PCV: 1 dose PCV at least 8 weeks after the most recent dose of PPSV23

Sickle cell disease and other hemoglobinopathies; anatomic or functional asplenia; congenital or acquired immunodeficiency; HIV infection; chronic renal failure; nephrotic syndrome; malignant neoplasms; leukemias, lymphomas, Hodgkin disease, and other diseases associated with treatment with immunosuppressive drugs or radiation therapy; solid organ transplantation; multiple myeloma.

Age 2–5 years

• Any incomplete series with:

- 3 PCV doses: 1 dose PCV (at least 8 weeks after any prior PCV dose)

- Less than 3 PCV doses: 2 doses PCV (8 weeks after the most recent dose and administered 8 weeks apart)

• No history of PPSV23: 1 dose PPSV23 (at least 8 weeks after any prior PCV doses and a dose 2 of PPSV23 5 years later)

Age 6–18 years

• No history of either PCV or PPSV23: 1 dose PCV, 2 doses PPSV23 (dose 1 of PPSV23 administered 8 weeks after PCV and dose 2 of PPSV23 administered at least

Poliovirus vaccination

(minimum age: 6 weeks)

Routine vaccination

• 4-dose series at ages 2, 4, 6–18 months, 4–6 years; administer the final dose on or after age 4 years and at least 6 months after the previous dose.

• 4 or more doses of IPV can be administered before age 4 years when a combination vaccine containing IPV is used. However, a dose is still recommended on or after age 4 years and at least 6 months after the previous dose.

Catch-up vaccination

• In the first 6 months of life, use minimum ages and intervals only for travel to a polio-endemic region or during an outbreak.

• IPV is not routinely recommended for U.S. residents age 18 years or older.

Series containing oral polio vaccine (OPV), either mixed OPV-IPV or OPV-only series:

• Total number of doses needed to complete the series is the same as that recommended for the U.S. IPV schedule. See www.cdc.gov/mmwr/volumes/66/wr/mm6601a6.htm?s_cid=mm6601a6_w.

• Only trivalent OPV (tOPV) counts toward the U.S. vaccination requirements.

- Doses of OPV administered before April 1, 2016, should be counted (unless specifically noted as administered during a campaign).

- Doses of OPV administered on or after April 1, 2016, should not be counted.

- For guidance to assess doses documented as "OPV," see www.cdc.gov/mmwr/volumes/66/wr/mm6606a7.htm?s_cid=mm6606a7_w.

• For other catch-up guidance, see Table 2.

Special situations

• Adolescents aged 18 years at increased risk of exposure to poliovirus with:

- No evidence of a complete polio vaccination series (i.e., at least 3 doses): administer remaining doses (1, 2, or 3 doses) to complete a 3-dose series

- Evidence of completed polio vaccination series (i.e., at least 3 doses): may administer one lifetime IPV booster

For detailed information, see: www.cdc.gov/vaccines/vpd/polio/hcp/recommendations.html

Special situations

- Adolescents aged 18 years at increased risk of exposure to poliovirus with:
 - No evidence of a complete polio vaccination series (i.e., at least 3 doses): administer remaining doses (1, 2, or 3 doses) to complete a 3-dose series
 - Evidence of completed polio vaccination series (i.e., at least 3 doses): may administer one lifetime IPV booster

For detailed information, see:

www.cdc.gov/vaccines/vpd/polio/hcp/recommendations.html

Appendix

Contraindications and Precautions

Table 1 Recommended Adult Immunization Schedule by Age Group, United States, 2023

Vaccine	19–26 years	27–49 years	50–64 years	≥65 years
COVID-19	2- or 3- dose primary series and booster (See Notes)			
Influenza inactivated (IIV4) or Influenza recombinant (RIV4)	1 dose annually			
Influenza live, attenuated (LAIV4)	1 dose annually			
Tetanus, diphtheria, pertussis (Tdap or Td)	1 dose Tdap each pregnancy; 1 dose Td/Tdap for wound management (see notes)			
	1 dose Tdap, then Td or Tdap booster every 10 years			
Measles, mumps, rubella (MMR)	1 or 2 doses depending on indication (if born in 1957 or later)			For healthcare personnel, see notes
Varicella (VAR)	2 doses (if born in 1980 or later)	2 doses		
Zoster recombinant (RZV)	2 doses for immunocompromising conditions (see notes)		2 doses	
Human papillomavirus (HPV)	2 or 3 doses depending on age at initial vaccination or condition	27 through 45 years		
Pneumococcal (PCV15, PCV20, PPSV23)	1 dose PCV15 followed by PPSV23 OR 1 dose PCV20 (see notes)			1 dose PCV15 followed by PPSV23 OR 1 dose PCV20
Hepatitis A				

Table 1 Recommended Adult Immunization Schedule by Age Group, United States, 2023

Vaccine	19–26 years	27–49 years	50–64 years	≥65 years
COVID-19	2- or 3- dose primary series and booster (See Notes)			
Influenza inactivated (IIV4) or Influenza recombinant (RIV4)	1 dose annually			
Influenza live, attenuated (LAIV4)	1 dose annually			
Tetanus, diphtheria, pertussis (Tdap or Td)	1 dose Tdap each pregnancy; 1 dose Td/Tdap for wound management (see notes)			
	1 dose Tdap, then Td or Tdap booster every 10 years			
Measles, mumps, rubella (MMR)	1 or 2 doses depending on indication (if born in 1957 or later)			For healthcare personnel, see notes
Varicella (VAR)	2 doses (if born in 1980 or later)		2 doses	
Zoster recombinant (RZV)	2 doses for immunocompromising conditions (see notes)		2 doses	
Human papillomavirus (HPV)	2 or 3 doses depending on age at initial vaccination or condition	27 through 45 years		
Pneumococcal (PCV15, PCV20, PPSV23)	1 dose PCV15 followed by PPSV23 OR 1 dose PCV20 (see notes)			1 dose PCV15 followed by PPSV23 OR 1 dose PCV20
Hepatitis A (HepA)	2, 3, or 4 doses depending on vaccine			
Hepatitis B (HepB)	2, 3, or 4 doses depending on vaccine or condition			
Meningococcal A, C, W, Y (MenACWY)	1 or 2 doses depending on indication, see notes for booster recommendations			
Meningococcal B (MenB)	2 or 3 doses depending on vaccine and indication, see notes for booster recommendations			

Table 1 Recommended Adult Immunization Schedule by Age Group, United States, 2023

Vaccine	19–26 years	27–49 years	50–64 years	≥65 years
COVID-19	2- or 3- dose primary series and booster (See Notes)			
Influenza inactivated (IIV4) or Influenza recombinant (RIV4)	1 dose annually			
Influenza live, attenuated (LAIV4)	1 dose annually			
Tetanus, diphtheria, pertussis (Tdap or Td)	1 dose Tdap each pregnancy; 1 dose Td/Tdap for wound management (see notes)			
	1 dose Tdap, then Td or Tdap booster every 10 years			
Measles, mumps, rubella (MMR)	1 or 2 doses depending on indication (if born in 1957 or later)			For healthcare personnel, see notes
Varicella (VAR)	2 doses (if born in 1980 or later)		2 doses	
Zoster recombinant (RZV)	2 doses for immunocompromising conditions (see notes)		2 doses	
Human papillomavirus (HPV)	2 or 3 doses depending on age at initial vaccination or condition	27 through 45 years		
Pneumococcal (PCV15, PCV20, PPSV23)	1 dose PCV15 followed by PPSV23 OR 1 dose PCV20 (see notes)			1 dose PCV15 followed by PPSV23 OR 1 dose PCV20
Hepatitis A (HepA)	2, 3, or 4 doses depending on vaccine			
Hepatitis B (HepB)	2, 3, or 4 doses depending on vaccine or condition			
Meningococcal A, C, W, Y (MenACWY)	1 or 2 doses depending on indication, see notes for booster recommendations			
Meningococcal B (MenB)	2 or 3 doses depending on vaccine and indication, see notes for booster recommendations			
	19 through 23 years			
Haemophilus influenzae type b (Hib)	1 or 3 doses depending on indication			
<div><div>Recommended vaccination for adults who meet age requirement, lack documentation of vaccination, or lack evidence of past infection</div><div>Recommended vaccination for adults with an additional risk factor or another indication</div><div>Recommended vaccination based on shared clinical decision-making</div><div>No recommendation/Not applicable</div></div>				

Table 2

The Medical Indications Table